1892.]

NOTES ON TRICHOPTERA AND NEUROPTERA FROM IRELAND. BY KENNETH J. MORTON.

During the first half of last August I collected in the neighbour-hood of Glasslough in Co. Monaghan, Ireland, and I now record those species of *Trichoptera* which have not previously been taken in the locality, and also the dragon-flies and a few other *Neuroptera* which were observed. The weather was, on the whole, fine, and many caddisflies were abundant; dragon-flies were also about in great numbers, but other families, such as *Chrysopidæ* and *Hemerobiidæ*, were hardly represented at all.

The Trichoptera new to the district are as follows:—Limnophilus flavicornis, Fab.; L. xanthodes, McLach., 3 \(\cop ; L. lunatus, Curt.; L. auricula, Curt.; L. hirsutus, Pict., 1 \(\cop ; L. sparsus, Curt.; Leptocerus dissimilis, Ste.; Mystacides nigra, L.; M. azurea, L.; Ecetis ochracea, Curt.; Psychomyia pusilla, Fab.; Agraylea multipunctata, Curt.; Hydroptila femoralis, Eat.; H. forcipata, Eat.; and Ithytrichia lamellaris, Eat. Other interesting species, such as Agrypnia Pagetana, Curt., Glyphotælius pellucidus, Retz., Limnophilus decipiens, Kol., Leptocerus fulvus, Ramb., Ecetis furva, Ramb., Tinodes maculicornis, Pict., &c., were also taken.

Of dragon-flies Sympetrum scoticum, Don., swarmed on a bog; S. striolatum, Charp., was more generally distributed, but not so numerous. Libellula quadrimaculata, L., flew sparingly over the bog. Æschna juncea, L., common. Æ. grandis, L.: this fine insect was even commoner than juncea, and many examples of it were taken; while it came more conspicuously under notice during the hours of bright sunshine, when it might be seen chasing and capturing such large game as Charæas graminis and Hydræcia nictitans, it also followed its well known crepuscular habit, and one dull, warm evening I watched some examples carrying on their feeding operations along the shores of the lake until it was nearly dark. Lestes sponsa, Hans., this species was exceedingly common at both bog and lakes. Ischnura elegans, V. de Lind., also in great abundance. Enallagma cyathigerum, Charp., common. Another dragon-fly seen but not taken, was probably Cordulegaster annulatus.

Very few species belonging to the other Families of Neuroptera were collected. Amongst these may be mentioned Sisyra fuscata, Fab., which was swarming; and Psocus longicornis, Fab., and P. nebulosus, Ste.